

## Literatura:

- Daniel, M., Černý, V., Dusbábek, F., Honzáková, E., Olejníček, J. (1976): Influence of microclimate on the life cycle of the common tick *Ixodes ricinus* in thermophilic oak forest. *Folia Parasitologica*, 23: 327-342.
- Daniel, M., Černý, V., Dusbábek, F., Honzáková, E., Olejníček, J. (1977): Influence of microclimate on the life cycle of the common tick *Ixodes ricinus* in an open area in
- Daniel, M., Kolář, J. (1990): Using satellite data to forecast the occurrence of the common tick *Ixodes ricinus* (L.). *Journal of Hygiene, Epidemiology and Immunology*, 34: 243-252. comparison with forest habitats. *Folia Parasitologica*, 24: 149-160.
- Daniel, M., Kolář, J. (1993): Remote sensing recognition of *Ixodes ricinus* habitats using LANDSAT MSS imagery. *Mitteilungen Österreichischen Gesellschaft Tropenmedizin Parasitologie*, 15: 157-161.
- Daniel, M., Dusbábek, F. (1994): Micrometeorological and microhabitat factors affecting maintenance and dissemination of tick-borne diseases in the environment. In: Sonenshine, D. E., Mather, T. N. (Eds.), *Ecological Dynamics of Tick-borne Zoonoses*. 1994, New York – Oxford, Oxford University Press, 91-138.
- Daniel, M., Kolář, J., Zeman, P., Pavelka, K., Sádlo, J. (1998a): Predictive map of *Ixodes ricinus* high-incidence habitats and a tick-borne encephalitis risk assessment using satellite data. *Experimental and Applied Acarology*, 22: 417-433.
- Daniel, M., Kolář, J., Zeman, P., Pavelka, K., Sádlo, J. (1998b): Predikce míst zvýšeného rizika napadení klíštětem *Ixodes ricinus* a nákazy klíšťovou encefalitidou ve středočeském regionu na základě družicových dat. *Epidemiologie, mikrobiologie, imunologie*, 47: 3-11.
- Daniel, M., Kolář, J., Zeman, P., Pavelka, K., Sádlo, J. (1999): Tick-borne encephalitis and Lyme borreliosis: comparison of habitat risk assessment using satellite data. *Central European Journal of Public Health*, 7: 35-39.
- Daniel, M., Kříž, B. (2002): Klíšťová encefalitida v České republice.  
I. Predikční mapy zvýšeného výskytu klíštěte obecného *Ixodes ricinus* a možného rizika nákazy virem klíšťové encefalitidy v českých regionech (Daniel M., Kolář J., Beneš Č., Danielová V.).  
II. Mapy případů onemocnění klíšťovou encefalitidou registrovaných na území České republiky v letech 1971 – 2000 (Pejčoch M., Beneš Č., Vymazal J.). Project Climate Change and Adaptation Strategies for Human Health in Europe, EVK2-2000-00670. Státní zdravotní ústav, Praha, 2002.
- Daniel M., Danielová, V., Kříž, B., Jirsa, A., Nožička, J. (2003) Shift of the tick *Ixodes ricinus* and tick-borne encephalitis to higher altitudes in Central Europe. *Eur. J. Clin. Microbiol. Infect. Dis.*, 22: 327-328.
- Daniel, M., Danielová, V., Kříž, B., Beneš, Č. (2006): Tick-borne encephalitis. In: Menne, B., Ebi, K.L. (Eds). *Climate change and adaptation strategies for human health 1*, pp. 89- 205. Steinkopff, Springer, Darmstadt, 2006.
- Daniel, M., Kolář, J., Zeman, P. (2008a): Analysing and predicting the occurrence of ticks and tick-borne diseases using GIS. In: Bowman, A. S., Nuttall, P. A. (Eds.): *Ticks: biology, disease, and Control.*, pp. 377-407. Cambridge University Press, Cambridge, 2008.
- Daniel, M., Kříž, B., Danielová, V., Valter, J., Kott, I. (2008b): Correlation between meteorological factors and tick-borne encephalitis in the Czech Republic. *Parasitology Research*, (Suppl. 1) 103: S97-S107.
- Daniel, M., Kříž, B., Valter, J., Kott, I., Danielová, V., (2008c): The influence of meteorological conditions of the preceding winter on the incidence of the tick-borne

- encephalitis and Lyme borreliosis in the Czech Republic. *International Journal of Medical Mikrobiology*, 298, S1: 60-67.
- Daniel, M., Kříž, B., Danielová, V., Beneš, Č. (2008d): Sudden increase in tick-borne encephalitis cases in the Czech Republic, 2006. *Int. J. Med. Microbiol.*, 298, S 1: 81-87.
- Daniel, M., Materna, J., Hönl V., Metelka L., Danielová V., Harčarik J., Kliegrová S., Grubhoffer L. (2009): Vertical distribution of the tick *Ixodes ricinus* and tick-borne Pathogens in the Northern Moravian mountains correlated with climate warming (Jeseníky Mts., Czech Republic). *Central European Journal of Public Health*, 17: 139-145.
- Danielová, V., Beneš, Č. (1997): Zvýšený výskyt klíšťové encefalitidy v České republice. *Praktický lékař*, 77: 580-583.
- Danielová, V., Kříž, B., Daniel, M., Beneš, Č., Valter, J., Kott, I. (2004): Vliv změn klimatu na výskyt klíšťové encefalitidy v České republice v uplynulých dvaceti letech. *Epidemiologie, mikrobiologie, imunologie*, 53: 174-181.
- Danielová V., Rudenko, N., Daniel, M., Holubová, J., Materna, J., Golovchenko, M., Schwarzová, L. (2006): Extension of *Ixodes ricinus* ticks and agents of tick-borne diseases to mountain areas in the Czech Republic. *Int. J. Med. Microbiol.* 296, S1, 48-53
- Danielová, V., Schwarzová, L., Materna, J. Daniel, M., Metelka, L., Holubová, J., Kříž, B. (2008a). Tick-borne encephalitis virus expansion to higher altitudes correlated with climate warming. *Int. J. Med. Microbiol.* 298, S1, 68 - 72.
- Danielová, V., Kliegrová S., Daniel, M., Beneš, Č. (2008b): Influence of climate warming on tick-borne encephalitis expansion to higher altitudes over last decade (1997-2006) in the Highland Region (Czech Republic). *Central European Journal of Public Health*, 16: 4-11.
- Danielová, V. , Daniel, M., Schwarzová, L., Materna, J. Rudenko, N., Golovchenko, M., Holubová, J., Grubhoffer, L., Kilián, P. (2009): Integration of a tick-borne encephalitis virus and *Borrelia burgdorferi* sensu lato into mountain ecosystems, following a shift in the altitudinal limit of distribution of their vector, *Ixodes ricinus* (Krkonoše Mts., Czech Republic). *Vector-borne and Zoonotic Diseases* 9 (4): v tisku.
- Kříž, B., Beneš, Č., Danielová, V., Daniel, M. (2004): Socio-economic conditions and other anthropogenic factors influencing tick-borne encephalitis incidence in the Czech Republic. *Int. J. Med. Microbiol.*, 293, Suppl. 37, 63-68.
- Materna, J., Daniel, M., Metelka, J., Harčarik, J. (2008): The vertical distribution, density and the development of the tick *Ixodes ricinus* in mountain areas influenced by climate changes (The Krkonoše Mts., Czech Republic). *International Journal of Medical Mikrobiology*, 298, S1: 25-37.
- Kříž B, Benes C, Daniel M.(2009), Alimentary transmission of tick-borne encephalitis in the Czech Republic (1997-2008), *Epidemiol Mikrobiol Imunol.* Apr;58(2):98-103.